

Altered Protein Expression in Peach (*Prunus persica*) Following Fruit Bagging

Wei Zhang¹, Xiaomeng Zhao^{2,4}, Mengya Shi³, Aizhen Yang^{2,4}, Baoguang Hua², and Yueping Liu^{2,4*}

¹College of Food Science and Engineering, Beijing University of Agriculture, Beinong Road 7, Huilongguan, Changping District, Beijing 102206, China

²College of Biological Science and Engineering, Beijing University of Agriculture, Beinong Road 7, Huilongguan, Changping District, Beijing 102206, China

³College of Plant Science and Technology, Beijing University of Agriculture, Beinong Road 7, Huilongguan, Changping District, Beijing, 102206 China

⁴Key Laboratory of Urban Agriculture (North) of Ministry of Agriculture P. R. China, Beijing University of Agriculture, Beijing 102206, China

*Corresponding author: liuyueping@bua.edu.cn

Abstract

Fruit bagging has been widely practiced in peach cultivation to produce high quality and unblemished fruit. Moreover, fruit bagging has been utilized to study the effect of shading on the quality of fruit. We conducted a proteomic analysis on peach fruit to elucidate the biochemical and physiological events that characterize the effect of bagging treatment. Comparative analysis of 2D electrophoresis (2-DE) gels showed that relative protein levels differed significantly at 125 DAFB (days after full bloom), as well as at 133 DAFB in fruit that had been bagged until 125 DAFB, followed by exposure to sunlight. Most of the proteins with altered expression were identified by MALDI TOF/TOF. Twenty-one proteins with differential expression among the groups were identified at 125 DAFB, while thirty proteins with differential expression among the groups were identified at 133 DAFB. The analysis revealed that expression of proteins involved in photosynthesis, stress responses, and biochemical processes influencing metabolism were altered during bagging treatment, suggesting that regulation of the synthesis of carbohydrates, amino acids, and proteins influenced fruit size, solid/acid ratio, and peel color. This work provides the first characterization of proteomic changes in peach in response to fruit bagging treatment. Identifying and tracking protein changes may allow us to better understand the mechanisms underlying the effects of bagging treatment.

Additional key words: 2D electrophoresis, bagging treatment, fruit development, MALDI TOF/TOF, proteomics

Korean J. Hortic. Sci. Technol. 34(1):32-45, 2016
http://dx.doi.org/10.12972/kjhst.20160019

pISSN : 1226-8763
eISSN : 2465-8588

Received: July 15, 2015

Revised: October 20, 2015

Accepted: January 25, 2016

Copyright©2016 Korean Society for Horticultural Science.

Introduction

Peach (*Prunus persica*) is among the most economically important fruit producing rosaceous crops and a model species used in studies of stone fruit in the *Rosaceae* family (Shulaev et al., 2008). Fruit bagging while on the tree is extensively practiced in China to improve fruit quality via promotion of anthocyanin synthesis. Upon removal of the bag from the treated plants, the pigmentation of peach,

apple, pear, and mango fruit is improved (Liu et al., 2013; Yu et al., 2012; Jia et al., 2005; Hofman et al., 1997). In addition to improving color, the practice of bagging prevents infestation by insect pests, reduces egg laying by female flies, inhibits disease, reduces physical damage (scars and scratches) (Amarante et al., 2002), and inhibits splitting and sunburn (Bentley and Viveros, 1992), while avoiding direct contact with pesticides and thus greatly reducing agrochemical residues. Although laborious, the practice of bagging is safer and easier than pesticide application, while providing farmers with more reliable estimates of projected harvests than are possible with non-bagging cultivation methods. Finally, bagging increases marketable fruit yield by reducing cosmetic defects.

Studies of the physiological mechanisms underlying the effects of bagging have primarily focused on anthocyanin synthesis following bagging of apple (Liu et al., 2013) and pear (Yu et al., 2012) trees. It is believed that bagging increases the sensitivity of fruit to light when the bag is removed, thus enhancing light-stimulated anthocyanin synthesis (although anthocyanin production in apple fruit also requires cool night temperatures). In addition, bagging may increase the abundance of volatile aromatic compounds in the skin of peach fruit, which may improve flavor (Jia et al., 2005). A recent study demonstrated that accumulation of *n*-hexanal and (*E*)-2-hexenal, which contribute to the characteristic aroma of peach fruit, is related to the particular sunlight transmission properties of the bags used to cover the fruit (Shen et al., 2014). In addition to color and aroma, fruit bagging also affects qualities of the flesh of the fruit, including sweetness and acidity. Most studies have reported that fruit bagging led to reduced contents of soluble carbohydrates, phenolic compounds, and organic acids (Lima et al., 2013) or had no changes in soluble carbohydrate content. Reports have mainly focused on the effects of fruit bagging on general fruit quality; however, fruit quality is the result of the combined action of numerous signaling and metabolic pathways during fruit development. Therefore, it is necessary to identify and quantify early responses to bagging at the protein level.

Recently, proteomics, the systematic study of global changes in protein expression, has been used in studies of peach fruit development (Hu et al., 2011; Prinsi et al., 2011; Nilo et al., 2010), post-harvest treatment (Zhang et al., 2011), and heat stress (Liu et al., 2012). Proteomics studies of peach fruit have provided insight into the physiological mechanisms underlying peach development. Feng et al. (2011) reported that fruit bagging altered the expression levels of numerous proteins in pear fruit.

The purpose of this study was to utilize proteomics to measure changes in protein expression following bagging of peach fruit, with the primary goals of identifying new biomarkers of fruit quality and evaluating alterations in adaptive responses in fruit, to provide a detailed characterization of the effect of fruit bagging on fruit quality at the proteomic level.

Materials and Methods

Plant Material and Growth Conditions

Peach (*P. persica* L. cv. 'No. 24') fruit were grown in an experimental peach orchard at the Beijing University of Agriculture (Pinggu District, Beijing, China) using conventional practices. Three neighboring peach trees with similar loads were the source of fruit. Bagging with two layers of pale-black fruit bags (Xintaiguodai factory, Shangdong, China) was carried out 50 days after full bloom (DAFB). Half of the fruit on each tree was bagged. Bagged and non-bagged fruit samples with no symptoms of pests and diseases were harvested at 125 DAFB from three different trees. On the same day, some of the bagged fruit was exposed to sunlight; this fruit was harvested at 133 DAFB. Non-bagged fruit was also

harvested at 133 DAFB. For each treatment and control group, 21 fruits were harvested from three trees at random to avoid biasing the samples, after which the fruits were randomly divided into three groups, which served as three biological replicates ($n = 7$ in each replicate). The mesocarps from individual fruit in each treatment group in each replicate were pooled, immediately frozen in liquid nitrogen, and stored at -80°C . For determination of fruit quality, 10 fruits were harvested from each tree.

Fruit Quality Trait Determination

The fresh weight (FW), vertical diameter (VD), and transverse diameter (TD) were the averages of the measurements from 10 fruits. A TA-XT2i plus texture analyzer (Stable Micro System, Godalming, England) equipped with a 7.9-mm diameter head was applied for fruit firmness measurement as described by Zhang et al. (2010).

Skin color was visually assessed as the percentage of skin area showing red color and ranked on a scale of 0 (no red color) to 10 (intense red). Anthocyanin was extracted from mesocarp taken from the reddest part of each fruit with 25 mL of 1% HCl in methanol. The absorbance at 528 nm was measured by a spectrophotometer using the method reported by Chaovanalikit (2004). For the chlorophyll content assay, circular pieces were cut from the mesocarp and extracted with methanol using a mortar, after which the extracted pigments were centrifuged at $8000 \times g$ (Sigma 3K-30, Germany) for 5 min to make the extract fully transparent. The resulting extracts were immediately assayed spectrophotometrically (SmartSpec Plus, Bio-Rad, USA). The specific absorption coefficients of Chl *a* and Chl *b* were taken from Lichtenthaler (1987).

Measurements of soluble solids content (SSC) and titratable acidity (TA) were conducted on juice samples collected from each treatment group. The SSC in opposite parts of each fruit was measured with a refractometer (Atago CM780N, Japan). TA was determined by titration to pH 8.1 with 0.05 N NaOH. The TA results were expressed in grams of malic acid per liter of juice.

Protein Extraction and Quantification

Total protein was extracted and purified from ground fruit mesocarp via phenol extraction followed by ammonium acetate methanol precipitation, as described by Faurobert et al. (2007) and Hu et al. (2011). The protein concentration in each extract was determined using the Non-Interfering Protein Assay TM Kit (EMD Biosciences, Germany) to remove interfering compounds, with bovine serum albumin (BSA) as the standard for the calibration curve. The re-solubilized protein samples were stored at -80°C prior to use.

2-D Electrophoresis

Three gels were run for each extract. Briefly, IPG strips (pH 4-7, 17 cm, ReadyStrip, Bio-Rad, Hercules, CA, USA) were actively rehydrated (50 V) overnight with 300 μL of IEF buffer containing 800 μg of total protein. Isoelectric focusing (IEF), equilibrating, and 2-D electrophoresis (2-DE) were performed according to the methods of Hu et al. (2011). 2-DE gels were stained with Coomassie Brilliant Blue-G250 (CBB) following a reported blue-silver protocol (Candiano et al., 2004).

Image Acquisition and Analysis

Gel images were acquired using a UMAX 2100 XL Scanner (UMAX Technologies, Dallas, TX, USA) and analyzed with PDQuest Advanced software version 8.0.1 (Bio-Rad, Hercules, CA, USA). Spot detection and matching were performed automatically, followed by manual inspection to correct any error prior to final data analysis. After normalization of the protein spot volume against the spot volume of the entire gel, the percentage volume of each spot was averaged for the three gels. Statistically significant changes in protein abundance were determined using three sequential data analysis criteria. First, a protein spot had to be present in all replicated gels. Next, a two-fold change in normalized spot volume was considered as indicative of a significant quantitative variation. Third, p -values < 0.05 were considered to be statistically significant by Duncan's multiple range test.

MS (mass spectrometry) Analysis and Protein Identification

Spots detectable in three replicates were excised and digested with trypsin according to the procedure of Hu et al. (2011). The extracted trypsin fragments were analyzed by 4800 MALDITOF/TOF™ (AB Sciex, US) according to the manufacturer's protocol. The raw spectra were manually filtered by PEAKERAZOR to detect potential contaminants and submitted to MASCOT software to search the NCBIInr and SwissProt databases.

Statistical Analysis

All statistical analyses were performed using SPSS 14.0 (SPSS Inc., Chicago, IL, USA). Data were analyzed using analysis of variance (ANOVA), after which group means were compared by Duncan's multiple range test (DMRT) with a significance threshold of $p < 0.05$.

Results and Discussion

Effect of Bagging on Fruit Quality

Differences in the external morphological characteristics of bagged and non-bagged peach fruit are shown in Fig. 1. At 125 DAFB, the ground color of bagged fruit was slightly yellow (Fig. 1B), while the skin of the non-bagged fruit was green (Fig. 1A). The bagged fruit rapidly turned red following bag removal (Fig. 1D), at which point the skin of fruit had better market acceptance than that of the non-bagged fruit (Fig. 1C). Bagging treatment increased the FW and size of peach fruit, but there was no difference in fruit firmness after bagging treatment (Table 1).

Table 1. Effect of bagging on FW, size and firmness in peach fruit at two developmental stages.

Treatment	Control		Bagged	
	125 DAFB	133 DAFB	125 DAFB	133 DAFB
FW (g)	195.61 ^c	231.86 ^b	226.51 ^b	244.92 ^a
VD (cm)	6.29 ^c	7.29 ^b	7.12 ^b	7.83 ^a
TD (cm)	5.98 ^c	6.57 ^b	6.43 ^b	7.06 ^a
Firmness (N)	35.15 ^a	30.31 ^b	34.74 ^a	31.16 ^b

Means followed by different letters within a row were separated by LSD (5%). Each treatment had three replicates. FW: fresh weight; VD: vertical diameter; TD: transverse diameter.




Fig. 1. Differences in the external morphological characteristics of bagged and non-bagged peach fruit. Each panel consist of a top-view and side-view of the left and right peach fruit. (A) non-bagged fruit at 125 DAFB, (B) bagged fruit at 125 DAFB, (C) non-bagged fruit at 133 DAFB, (D) fruit bagged until 125 DAFB and exposed to sunlight until harvest and evaluation at 133 DAFB.

There was little difference in SSC content and no difference in TA content among the treatment groups. The solid/acid ratio of the bagged fruit at 125 DAFB and the debagged fruit at 133 DAFB were slightly lower than those of the non-bagged and bagged fruit at the same stages (Table 2). Furthermore, the solid/acid ratio increased with the developmental stage.

Skin color was significantly affected by bagging (Table 3). The percentage of skin area with a blush of color was larger in the debagged fruit than in the non-bagged fruit at 133 DAFB. Anthocyanin content was not significantly changed by

Table 2. Effect of bagging on SSC, TA and solid/acid ratio in peach fruit at two developmental stages.

Treatment	Control		Bagged	
	125 DAFB	133 DAFB	125 DAFB	133 DAFB
SSC (%)	12.88 ^{ab}	14.13 ^a	11.97 ^b	13.02 ^{ab}
TA (%)	0.20 ^a	0.19 ^a	0.22 ^a	0.20 ^a
Solid/acid ratio	64.40 ^b	74.37 ^a	54.41 ^c	65.10 ^b

Means followed by different letters within a row were separated by LSD (5%). Each treatment had three replicates.

Table 3. Effect of bagging on peach fruit pigmentation at two developmental stages

Treatment	Control		Bagged	
	125 DAFB	133 DAFB	125 DAFB	133 DAFB
Color intensity ^z (%)	19.88 ^c	64.81 ^b	5.97 ^d	73.40 ^a
Anthocyanin content (OD528)	0.32 ^{ab}	0.44 ^a	0.21 ^b	0.50 ^a
Chl <i>a</i> (mg · 100 g ⁻¹ FW)	0.41 ^a	0.47 ^a	0.18 ^b	0.25 ^b
Chl <i>b</i> (mg · 100 g ⁻¹ FW)	0.91 ^a	0.74 ^a	0.40 ^b	0.37 ^b
Chl (<i>a</i> + <i>b</i>) (mg · 100 g ⁻¹ FW)	1.32 ^a	1.21 ^a	0.58 ^b	0.62 ^b

Means followed by different letters within a row were separated by LSD (5%). Each treatment had three replicates.

^zColoration ratio on the equator

bagging treatment at 125 DAFB, whereas the fruit re-exposed to sunlight showed increased anthocyanin content at 133 DAFB (Table 3). In a previous study, fruit covered with bags appeared bright red and had high L values, which accounted for their good visual quality (Jia et al., 2005). The chlorophyll content of the non-bagged fruit was about twice that of the bagged fruit at 125 DAFB. In the fruit re-exposed to sunlight, the chlorophyll content in the debagged fruit was lower than that of the non-bagged fruit at 133 DAFB (Table 3).

Proteomic Profiles in Natural and Bagged Peach Fruit during Development

The proteomic profiles of the mesocarp from the bagged and non-bagged fruit were analyzed at 125 and 133 DAFB. Protein spots in all images were automatically detected and matched by PDQuest software, manually corrected spot-to-spot to identify various errors caused by noise and false spots, and subjected to analysis to calculate the effective number of spots (Fig. 2). Average proteomic maps showed approximately 500 spots for each stage. Based on the criteria described above, a total of 103 protein spots were subjected to mass spectrometry analysis, of which 68 spots showed a good quality MS signal and were subjected to a search in the NCBI nr and SwissProt databases. Protein spots were analyzed by comparing their theoretical molecular weight (Mr.) and isoelectric point (pI) with experimental Mr. and pI as determined in 2-D gels. Twenty-one spots showed different expression levels in bagged and non-bagged fruit at 125 DAFB, whereas 30 spots showed different expression levels in bagged and non-bagged fruit at 133 DAFB, including up- and down-regulated




Fig. 2. Two-dimensional electrophoresis maps of total proteins from non-bagged fruit (A) and bagged fruit (B) at 125 DAFB, as well as non-bagged fruit (C) and bagged fruit exposed to sunlight (D) from 125 DAFB to 133 DAFB. The differentially expressed proteins are numbered. Proteins were separated over the pI range of 4-7 in the first dimension and on 12% SDS-polyacrylamide gels in the second dimension. Gels were stained with colloidal Coomassie G250.

proteins, as well as proteins present in some samples and not in others.

Protein Identification and Functional Analysis

The identified proteins and their functions are listed in Table 4. The differentially expressed proteins in bagged and non-bagged fruit at 125 DAFB were classified into the following functional categories: stress response and defense (47.6%), photosynthesis (4.8%), carbohydrate metabolism (9.5%), amino acid metabolism (9.5%), protein synthesis and degradation (9.5%), cytoskeleton and signaling transduction (4.8%), fatty acid metabolism (4.8%), aroma synthesis (4.8%) and unknown protein (4.8%) (Fig. 3A). The differentially expressed proteins in bagged and non-bagged fruit at 133 DAFB were involved in stress response and defense (43.3%), photosynthesis (13.3%), carbohydrate metabolism (13.3%), amino acid metabolism (16.7%), protein synthesis and degradation (3.3%), cytoskeleton and signaling transduction (6.7%), and flavonoid biosynthesis (3.3%) (Fig. 3B). Stress/defense proteins were the most represented group, followed by components associated with photosynthesis.




Fig. 3. Functional distribution of differentially expressed protein spots in bagged fruit at 125 DAFB (A) and in debagged fruit at 133 DAFB (B). Identified protein species were categorized into different classes according to the KEGG and MIPS databases. Percentual discrepancies with respect to categorization present were due to the eventual attribution of individual protein entries into multiple functional classes. Stress/defense proteins were the most represented group, followed by components associated with photosynthesis.

Table 4. Protein species showing quantitative changes after bagging as identified by 2-DE and MS procedures.

Spot ID	Protein name	Accession No.	Organism	Mascot score	Coverage (P) ^y	Theor. Mr./pI	Exp. Mr./pI	ratio (bagged/non-bagged)	
								125 DAFB	133 DAFB
Stress response and defense									
1819	Hsc70	AAB42159	<i>Lycopersicon esculentum</i>	103	0%(14)	71.47/5.18	70.00/5.4	0.26	Und.
1709	heat shock 70 kDa protein	P11143	<i>Zea mays</i>	103	25%(18)	70.87/5.22	76.40/5.5	Und.	0.49
6014	18.1 kDa class I heat shock	P19243	<i>esculentum</i>	67	0%(7)	18.07/5.83	19.00/6.5	2.44	4.79
6007	17.7 kDa heat shock protein	AAB63311	<i>Helianthus annuus</i>	84	10%(5)	17.66/6.19	19.00/6.55	**	**
4008	small heat shock protein	AAR99375	<i>Prunus persica</i>	161	34%(3)	17.38/5.98	17.00/6.1	25.75	5.48
2008	cytosolic class II low molecular weight heat shock protein	AF159562_1	<i>Prunus dulcis</i>	68	19%(2)	17.55/5.58	16.00/5.7	10.44	4.81
3206	abscisic stress ripening-like protein	AAL26889	<i>Prunus persica</i>	132	16%(8)	20.75/5.68	33.00/5.9	4.2	Und.
3202	abscisic stress ripening-like protein	AAL26889	<i>Prunus persica</i>	61	11%(3)	20.75/5.68	31.00/5.9	4.05	4.99
4202	abscisic stress ripening-like protein	AAL26889	<i>Prunus persica</i>	106	31%(4)	20.75/5.68	30.00/6.0	3.02	2.24
3203	abscisic stress ripening protein homolog	AAB97140	<i>Prunus armeniaca</i>	45	5%(1)	21.23/5.64	31.00/5.9	Und.	3.79
6716	polyphenol oxidase precursor	AAC28935	<i>Prunus armeniaca</i>	62	4%(2)	67.44/6.39	60.00/6.45	2.27	Und.
5711	polyphenol oxidase precursor	AAC28935	<i>Prunus armeniaca</i>	271	7%(4)	67.44/6.39	61.00/6.5	Und.	2.53
5705	polyphenol oxidase precursor	AAC28935	<i>Prunus armeniaca</i>	56	3%(2)	67.44/6.39	63.00/6.3	Und.	2.37
5716	polyphenol oxidase precursor	AAC28935	<i>Prunus armeniaca</i>	118	7%(4)	67.44/6.39	66.00/6.4	Und.	3.03
6702	polyphenol oxidase precursor	AAC28935	<i>Prunus armeniaca</i>	49	4%(2)	67.44/6.39	61.00/6.4	Und.	2.32
3011	thioredoxin H	AAL26915	<i>Prunus persica</i>	48	19%(2)	14.66/5.62	18.30/5.9	0.29	0.38
Photosynthesis									
1218	oxygen-evolving enhancer protein 1	Q40459	<i>Nicotiana tabacum</i>	87	5%(9)	35.21/5.89	29.00/5.6	*	*
3209	oxygen-evolving enhancer protein 1, chloroplastic	O49079	<i>Fritillaria agrestis</i>	37	5%(1)	35.08/6.26	37.00/5.9	Und.	0.07
5401	delta-aminolevulinic acid dehydratase, chloroplastic	P30124	<i>Pisum sativum</i>	110	10%(3)	44.02/5.67	41.00/6.25	Und.	2.59
7503	ribulose-1,5-bisphosphate carboxylase/oxygenase large subunit	AAB66923	<i>Combretocarpus rotundatus</i>	147	8%(4)	49.07/6.57	55.60/6.7	Und.	0.45
Carbohydrate metabolism									
6612	UTP-glucose-1-phosphate uridylyltransferase	O64459	<i>Pyrus pyrifolia</i>	52	22%(10)	51.98/5.99	55.60/6.5	2.33	2.27
5806	phosphoglucomutase, cytoplasmic	Q9SM60	<i>Pisum sativum</i>	49	2%(2)	63.51/5.46	66.40/6.3	0.41	Und.
1412	alpha-amylase	XP_003589186	<i>Medicago truncatula</i>	57	2%(1)	46.97/5.17	39.00/5.3	Und.	*
5709	malate dehydrogenase (oxaloacetate-decarboxylating)(NADP ⁺)	NP_179580	<i>Arabidopsis thaliana</i>	89	5%(3)	64.52/6.32	64.00/6.4	Und.	2.40
6808	malate dehydrogenase (oxaloacetate-decarboxylating)(NADP ⁺)	NP_179580	<i>Arabidopsis thaliana</i>	56	2%(2)	64.52/6.32	66.40/6.6	Und.	2.20
Amino acid metabolism									
5313	probable aldo-keto reductase 1	C6TBN2	<i>Glycine max</i>	66	10%(3)	38.46/6.14	38.60/6.5	0.42	Und.
5510	S-adenosyl-L-homocysteine hydrolase	CAI56440	<i>Cicer arietinum</i>	75	0%(12)	53.35/5.53	50.00/6.5	0.12	0.31
5404	S-adenosylmethionine synthetase 1	Q56TU4	<i>Daucus carota</i>	92	29%(15)	48.81/6.08	45.00/6.3	Und.	0.44
5403	S-adenosylmethionine synthetase-2	BAC81655	<i>Pisum sativum</i>	130	0%(8)	37.61/6.27	42.70/6.3	Und.	0.40
4309	arginase 2	NP_001233851	<i>Solanum lycopersicum</i>	78	6%(2)	36.94/5.60	31.00/6.0	Und.	0.40
5608	ketol-acid reductoisomerase	O82043	<i>Pisum sativum</i>	88	0%(9)	62.81/6.62	60.00/6.4	Und.	2.39
Protein synthesis and degradation									
0212	proteasome subunit alpha type-1-A	P34066	<i>Arabidopsis thaliana</i>	61	6%(1)	30.69/4.99	34.60/5.2	2.01	Und.
8411	ATP-dependent Clp protease proteolytic subunit 3, chloroplastic	Q9SXJ6	<i>Arabidopsis thaliana</i>	30	10%(6)	34.19/7.6	40.00/7.3	2.98	Und.
1302	DNA damage-inducible protein 1-like	XP_003532197	<i>Glycine max</i>	83	8%(2)	44.70/5.01	40.00/5.6	Und.	0.40

Spot ID	Protein name	Accession No.	Organism	Mascot score	Coverage (P) ^y	Theor. Mr./pI	Exp. Mr./pI	ratio (bagged/non-bagged)	
								125 DAFB	133 DAFB
Cytoskeleton and signaling transduction									
1607	tubulin alpha-1 chain	XP_002285721	<i>Vitis vinifera</i>	82	25%(11)	50.27/4.95	55.60/5.5	2.06	2.34
3403	actin-1	P23343	<i>Daucus carota</i>	49	7%(2)	42.25/5.64	42.00/5.9	Und.	3.76
Fatty acid metabolism									
5503	biotin carboxylase	AAK60339	<i>Brassica napus</i>	182	8%(4)	58.95/6.52	64.00/6.4	0.26	Und.
Aroma synthesis									
1602	mandelonitrile glucosyltransferase UGT85A19	ABV68925	<i>Prunus dulcis</i>	159	10%(6)	54.18/5.03	55.60/5.4	3.22	Und.
Flavonoid biosynthesis									
4304	leucoanthocyanidin dioxygenase	ABX89941	<i>Prunus persica</i>	92	0%(10)	40.45/5.46	39.00/6.1	Und.	2.52
Unknown protein									
2308	predicted protein	XP_002326572	<i>Populus trichocarpa</i>	92	0%(1)	45.00/6.11	38.00/5.9	*	Und.

Und.: undetectable.

*only observed in non-bagged fruit.

**only observed in bagged fruit.

^zIncreasing/decreasing index (fold-change) was calculated as the ratio of the relative expression level in the bagged group to that of the non-bagged group. Proteins were considered to be differentially expressed when the relative fold-change was > 2.0 or < 0.5. Functional grouping was performed according to the KEGG and MIPS databases.

^yAmino acid coverage (%) and the number of unique peptides (P).

Proteins Associated with Stress Response and Defense

Bagging subjects fruit to varying degrees of light deprivation and heat stress during the ripening process. Heat shock proteins (HSPs) can be induced by stimuli associated with bagging, including high temperature (Lara et al., 2009) and ripening (Nilo et al., 2010). Induction of small heat shock protein (sHSP) expression prevents protein misfolding under adverse conditions, including various stressors (Zou et al., 2009). In this study, four differentially expressed sHSPs (spots 6014, 6007, 4008, and 2008) were markedly induced following bagging and debagging treatment. The effects of stress and ripening caused by bagging were indistinguishable in our study.

The HSP70 family includes both heat-inducible and constitutively expressed members (D'Ambrosio, 2013). HSP70 protein levels increase upon exposure to various stressors (Ambrosia, 2013) and change in an isoform-specific fashion during fruit ripening (Huang et al., 2011). Giribaldi et al. (2007) reported that the expression of some HSPs decreased at the onset of ripening, consistent with a lower rate of protein synthesis and reduced protein yield. In our study, HSP70 (spot 1819 at 125 DAFB and spot 1709 at 133 DAFB) expression was down-regulated under bagging and debagging treatment. However, the stress caused by bagging generally increases HSP70 expression (Polenta et al., 2007); therefore, the decreased HSP70 expression observed in our study may be a result of changes in the ripening process caused by bagging treatment.

The abscisic stress ripening-like proteins belong to the abscisic acid/water deficit stress (ABA/WDS)-induced protein family, which consists of plant proteins induced by water deficit stress or ABA stress and ripening. Protein abscisic stress ripening-like proteins scavenge reactive oxygen species (ROS) in vitro (Kim et al., 2012). In this study, expression of abscisic stress ripening-like proteins (spots 3206, 3202, 4202, and 3203) was up-regulated in peach fruit by bagging treatment. Bagging generally caused earlier ripening, indicating that abscisic stress ripening-like proteins can serve as biomarkers of the peach ripening process. A previous study found that abscisic stress ripening-like proteins were differentially

regulated in ripe fruit and fruit stored at a cold temperature (Nilo et al., 2010); further studies must be performed to determine the function of these proteins in peach fruit.

Polyphenol oxidase (PPO) catalyzes the hydroxylation of monophenols to o-dihydroxyphenols and the oxidation of o-dihydroxyphenols to o-quinones (Cabanés et al., 2007). In most cases, PPO activity progressively increases as fruit matures, with the greatest increase in the final days before harvest. PPO activity increases 1.5 times after ethylene and respiratory peaks in Calanda peach (Ferrer et al., 2005). Expression of 5 PPO precursors (spots 6716, 5711, 5705, 5716, and 6702) was up-regulated by bagging at 125 DAFB and 133 DAFB. Thus, bagging treatment caused earlier fruit ripening, which was associated with PPO expression. These results suggest that PPO can serve as a biomarker of the peach fruit ripening process.

Thioredoxin H (spot 3011) has been associated with fruit ripening in several species, including peaches (Callahan et al., 1993; Manrique-Trujillo et al., 2007; Levi et al., 2006). In addition, thioredoxin H has been found to be responsive to abiotic stressors, including chilling injury and drought (Yan et al., 2006; Hajheidari et al., 2007). In this study, thioredoxin H expression was down-regulated by bagging at 125 DAFB and debagging at 133 DAFB. We were unable to distinguish whether the change in thioredoxin H expression associated with bagging treatment was the result of changes in the ripening process or stress.

Proteins Associated with Photosynthesis

Oxygen-evolving enhancer protein 1 and chloroplast oxygen-evolving enhancer protein 1 are components of photosystem II (PSII), which generates ATP to allow plant growth and development. In addition, PSII provides a precise scaffold for pigments and cofactors that mediate vectorial primary charge separation and electron transfer (Feng et al., 2011). Oxygen-evolving enhancer protein 1 (spot 1218) was only detected in non-bagged fruit, because such fruit have chlorophyll levels sufficient for photosynthetic activity. However, bagging treatment inhibited the accumulation and biosynthesis of chlorophyll. Even in fruit immediately exposed to sunlight after debagging, expression of chloroplast oxygen-evolving enhancer protein 1 (spot 3209) was markedly down-regulated. These results suggest that sunlight might be involved in the accumulation of phenolic compounds such as anthocyanin in peach peel.

Delta-aminolevulinic acid dehydratase catalyzes the synthesis of heme and Chl from the universal tetrapyrrole precursor (Shibata and Ochiai, 1977). Chl is the predominant tetrapyrrole in most plants. Delta-aminolevulinic acid dehydratase (spot 5401) expression was up-regulated in fruit that were re-exposed to sunlight after bagging. Chl is required for energy utilization when fruit are re-exposed to sunlight, leading to increased expression of delta-aminolevulinic acid dehydratase, which increases the photosynthetic capacity of the fruit.

Rubisco (spot 7503) is the key regulatory enzyme in the light-independent reactions of the Calvin cycle and catalyzes the first step in carbon fixation. The rate of photosynthesis was expected to decrease in bagged fruit subjected to low-light conditions. The bagged fruit at 125 DAFB and non-bagged fruit showed no difference in Rubisco expression. However, in comparison with non-bagged fruit, bagged fruit re-exposed to sunlight showed reduced Rubisco expression at 133 DAFB. The large change in photosynthesis occurring in fruit from 125 to 133 DAFB may have been a result of debagging.

These results show that expression levels of chloroplast oxygen-evolving enhancer protein 1, Rubisco, and delta-aminolevulinic acid dehydratase were changed in debagged fruit at 133 DAFB, indicating that fruit debagging affected biochemical pathways involved in photosynthetic carbon assimilation and protein turnover.

Proteins Associated with Carbohydrate Metabolism

Expression of UTP-glucose-1-phosphate uridylyltransferase (spot 6612), an enzyme involved in glycogenesis, was up-regulated by bagging. UTP-glucose-1-phosphate uridylyltransferase catalyzes the formation of UDP-glucose from glucose-1-phosphate and UTP. Bagging usually lowers the soluble carbohydrate content at harvest in apple (Watanabe et al., 2011) and pear fruit (Huang et al., 2009). Hiratsuka et al. (2012) reported that photosynthesis and phosphoenolpyruvate carboxylase (PEPC) activity were considerably inhibited after fruit bagging, while soluble carbohydrate content was reduced. In this study, the observed increase in UTP-glucose-1-phosphate uridylyltransferase expression was not associated with a change in soluble carbohydrate content. Other enzymes involved in soluble carbohydrate synthesis were not assessed in this study.

Malate dehydrogenase (MDH) reversibly catalyzes oxidation of malic acid to oxaloacetate via reduction of NAD^+ to NADH and can thus reduce cellular malate acid concentrations. MDH expression was increased in bagged fruit at 125 DAFB and in debagged fruit at 133 DAFB (spots 5709 and 6808), but TA did not differ significantly among the treatment groups (Table 2). The contents of intermediate products may be affected by multiple factors. The predominant organic compounds in ripe peach fruit are malic and citric acids, while quinic acid is present in smaller amounts (Moing et al., 1998); however, no enzyme capable of citric acid metabolism was identified as being regulated by bagging.

Phosphoglucomutase is involved in carbohydrate metabolism in peach fruit. In tomato fruit, phosphoglucomutase activity declines during early fruit development, as does the expression level of its cytosolic isoform. In tomato fruit, the highest level of phosphoglucomutase expression was found in ripe fruit, which was not the case for fruit of wild accessions (Kortstee et al., 2007). Phosphoglucomutase (spot 5806) expression was down-regulated in bagged fruit. Alpha-amylase (spot 1412) catalyzes starch degradation in peach fruit. Alpha-amylase was expressed in non-bagged fruit, but not in bagged fruit. The decreased protein levels of phosphoglucomutase and alpha-amylase in bagged peach fruit suggest that bagging may inhibit carbohydrate metabolism. Future studies will be aimed at identifying other enzymes involved in carbohydrate metabolism that may be regulated by fruit bagging.

Proteins Associated with Amino Acid Metabolism

The aldo-keto reductase (AKR, spot 5313) superfamily is a large enzyme group of NADP-dependent oxidoreductases with numerous roles in metabolism (Kanayama et al., 2014). S-adenosyl-L-homocysteine hydrolase (SAHH, spot 5510) catalyzes the reversible hydrolysis of S-adenosyl-L-homocysteine into adenosine and L-homocysteine, which can be regenerated into S-adenosyl-L-methionine (SAM) through L-methionine and sustains 1-aminocyclopropane-1-carboxylic acid (ACC) synthesis and methylation. S-adenosylmethionine synthetase (spot 5404 and 5403) catalyzes the rate-limiting step of the methionine cycle. SAM is a methyl donor in transmethylation reactions and the propylamino donor in polyamine biosynthesis. In plants, arginase (spot 4309) is involved in nitrogen remobilization following protein degradation, especially during seed germination (King and Gifford, 1997; Goldraij and Polacco, 2000). AKR, SAAH, and arginase expression levels were down-regulated following bagging treatment, suggesting that bagging inhibited nitrogen metabolism during ripening.

Ketol-acid reductoisomerase (KARI) is an oxidoreductase that acts on the CH-OH group of a donor molecule and utilizes NAD^+ or NADP^+ as an acceptor. KARI (spot 5608) expression was up-regulated at 133 DAFB.

Proteins Associated with Protein Synthesis and Degradation

Proteasome subunit alpha, type-1-A (spot 0212) is part of a multicatalytic proteinase complex characterized by its ability to cleave arginine, phenylalanine, tyrosine, leucine, and glutamic acid at neutral or slightly basic pH. Chloroplast ATP-dependent Clp protease proteolytic subunit 3 (spot 8411) is a Clp protease identified in *Arabidopsis thaliana* that is vital for chloroplast development and function (Sjögren et al., 2006). Removal of inactive polypeptides by proteases is particularly important during periods of stress, when the potential for protein damage is considerably increased. In bagged fruit, expression levels of proteasome subunit alpha, type-1-A and chloroplast ATP-dependent Clp protease proteolytic subunit 3 were up-regulated at 133 DAFB, suggesting that bagging altered protein metabolism.

DNA damage-inducible protein 1-like (spot 1302) acts as a linker between the 19S proteasome and polyubiquitinated proteins via UBA domains, facilitating degradation of ubiquitinated proteins. DNA damage-inducible protein 1-like expression was down-regulated at 133 DAFB, suggesting that proteasome-mediated degradation of ubiquitinated proteins may have been inhibited by bagging. Future studies will further evaluate changes in protein metabolism associated with fruit bagging.

Proteins Associated with Fatty Acid Metabolism, Aroma and Flavonoid Biosynthesis

Biotin carboxylase (spot 5503) participates in fatty acid biosynthesis. In fruit, fatty acids serve as precursors of aromatic compounds (Krammer et al., 1991). In bagged fruit, biotin carboxylase expression was down-regulated at 125 DAFB, suggesting that the reduced aroma of bagged fruit may have been a result of reduced precursor abundance.

A recent study suggested that mandelonitrile glucosyltransferase (spot 1602) GT85A19 activity is associated with bitterness in almond kernels (Franks et al., 2008). In this study, mandelonitrile glucosyltransferase expression was up-regulated in bagged fruit at 125 DAFB, which may have contributed to their reduced aroma. No further proteins with functions that could clearly be related to the aroma of peach fruit were identified as being differentially expressed in the bagged and non-bagged fruit.

Anthocyanidin synthase (spot 4304) participates in flavonoid biosynthesis. In this study, anthocyanidin synthase expression was up-regulated in bagged fruit at 133 DAFB, illustrating that bagging resulted in increased anthocyanin synthesis upon re-exposure of the bagged fruit to light.

Conclusion

Bagging has been used widely to improve fruit appearance, reduce pesticide residue levels, and increase commercial value. Fruit bagging is associated with changes in several characteristics of peach fruit, particularly fresh weight, skin color, and anthocyanin content. The present work reports a comprehensive proteomics analysis of peach fruit subjected to bagging in comparison with non-bagged fruit, with the goal of understanding the molecular mechanisms underlying the effects of fruit bagging. This investigation provides a preliminary overview of the important biological processes that contribute to the changes that occur in peach fruit following bagging. Functional classification of proteins with differential expression in bagged and non-bagged peach fruit revealed that the majority of regulated proteins were related to stress responses, photosynthesis, carbohydrate metabolism, amino acid metabolism, and synthesis of aromatic compounds. The functional

analysis of the proteins regulated by bagging was consistent with some of the fruit phenotypes produced by bagging. Proteome analysis can be used to link genotype and phenotype during plant growth and development, but few proteins directly relatable to phenotype were identified in this study due to the limited sensitivity of 2-D methods. These results enrich the body of information concerning the effect of bagging on protein metabolism in peach fruit. Future studies should endeavor to further explore the molecular mechanisms underlying the effects of bagging on fruit.

Acknowledgement: We gratefully acknowledge the National Science Foundation (31101526) and the Importation and Development of High-Caliber Talents Project of Beijing Municipal Institutions (CIT&TCD201404100) for funding this study.

Literature Cited

- Amarante, C., N.H. Banks, and S. Max. 2002. Preharvest bagging improves packout and fruit quality of pears (*Pyrus communis*). *N. Z. J. Crop Hortic.* 30:93-98.
- Bentley, W.J. and M. Viveros. 1992. Brown-bagging Granny Smith apples on trees stops codling moth damage. *California Agric.* 46:30-32.
- Cabanes, J., J. Escribano, F. Gandía-Herrero, F. García-Carmona, and M. Jiménez-Atiénzar. 2007. Partial purification of latent polyphenol oxidase from peach (*Prunus persica* L. cv. Catherina). Molecular properties and kinetic characterization of soluble and membrane-bound forms. *J. Agric. Food Chem.* 55:10446-10451.
- Callahan A.M., P.H. Morgens, and R.A. Cohen. 1993. Isolation and initial characterization of cDNAs for messenger-RNAs regulated during peach fruit-development. *J. Am. Soc. Hortic. Sci.* 118:531-537.
- Candiano, G., M. Bruschi, L. Musante, L. Santucci, G.M. Ghiggeri, B. Carnemolla, P. Orecchia, L. Zardi, and P.G. Righetti. 2004. Blue silver: a very sensitive colloidal Coomassie G-250 staining for proteome analysis. *Electrophoresis* 25:1327-1333.
- Chaovanalikit, A. and R.E. Wrolstad. 2004. Total anthocyanins and total phenolics of fresh and processed cherries and their antioxidant properties. *J. Food Sci.* 69:67-72.
- D'Ambrosio, C. S. Arena, M. Roccob, F. Verrillo, G. Novi, V. Viscosi, M. Marra, and A. Scaloni. 2013. Proteomic analysis of apricot fruit during ripening. *J. Proteomics* 78:39-57.
- Faurobert, M., C. Mihr, N. Bertin, T. Pawlowski, L. Negroni, N. Sommerer, and M. Causse. 2007. Major proteome variations associated with cherry tomato pericarp development and ripening. *Plant Physiol.* 143:1327-1346.
- Feng, S.Q., X.S. Chen, Y.M. Zhang, Y.L. Wang, Y. Song, X.C. Liu, X.G. Li, M. Li, J. Liu, Q.Z. Wang, and M.Y. Chen. 2011. Differential expression of proteins in red pear following fruit bagging treatment. *Protein J.* 30:194-200.
- Ferrer, A., S. Remon, A. Negueruela, and R. Oria. 2005. Changes during the ripening of the very late season Spanish peach cultivar Calanda - feasibility of using CIELAB coordinates as maturity indices. *Sci. Hortic.* 105:435-446.
- Franks, T.K., A. Yadollahi, M.G. Wirtensohn, J.R. Guerin, B.N. Kaoser, M. Sedgley, and C.M. Ford. 2008. A seed coat cyanohydrin glucosyltransferase is associated with bitterness in almond (*Prunus dulcis*) kernels. *Funct. Plant Biol.* 35:236-246.
- Giribaldi, M., I. Perugini, F.X. Sauvage, and A. Schubert. 2007. Analysis of protein changes during grape berry ripening by 2-DE and MALDI-TOF. *Proteomics* 7:3154-3170.
- Goldraj, A. and J.C. Polaco. 2000. Arginine degradation by arginase in mitochondria of soybean seedling cotyledons. *Planta* 210:652-658.
- Hajheidari, M., A. Eivazi, B.B. Buchanan, J.H. Wong, I. Majidi, and G.H. Salekdeh. 2007. Proteomics uncovers a role for redox in drought tolerance in wheat. *J. Proteome Res.* 6:1451-1460.
- Hiratsuka, S., Y. Yokoyama, H. Nishimura, T. Miyazaki, and K. Nada. 2012. Fruit photosynthesis and phosphoenolpyruvate carboxylase activity as affected by lightproof fruit bagging in Satsuma Mandarin. *J. Am. Soc. Hortic. Sci.* 137:215-220.
- Hofman, P.J., L.G. Smith, D.C. Joyce, G.I. Johnson, and G.F. Meiburg. 1997. Bagging of mango (*Mangifera indica* cv. 'Keitt') fruit influences fruit quality and mineral composition. *Postharvest Biol. Technol.* 12:83-91.
- Huang, C., F. Verrillo, G. Renzone, S. Arena, M. Rocco, A. Scaloni, and M. Marra. 2011. Response to biotic and oxidative stress in *Arabidopsis thaliana*: analysis of variably phosphorylated proteins. *J. Proteomics* 74:1934-1949.
- Huang, C.H., B. Yu, Y.W. Teng, J. Su, Q. Shu, Z.Q. Cheng, and L.Q. Zeng. 2009. Effects of fruit bagging on coloring and related physiology, and qualities of red Chinese sand pears during fruit maturation. *Sci. Hortic.* 121:149-158.
- Hu, H., Y. Liu, G.L. Shi, Y.P. Liu, R.J. Wu, A.Z. Yang, Y.M. Wang, B.G. Hua, and Y.N. Wang. 2011. Proteomic analysis of peach endocarp and mesocarp during early fruit development. *Physiol. Plant.* 142:390-406.
- Jia, H.J., A. Araki, and G. Okamoto. 2005. Influence of fruit bagging on aroma volatiles and skin coloration of 'Hakuho' peach (*Prunus persica* Batsch). *Postharvest Biol. Technol.* 35:61-68.

- Kanayama, Y., R. Mizutani, S. Yaguchi, A. Hojo, H. Ikeda, M. Nishiyama, and K. Kanahama. 2014. Characterization of an uncharacterized aldo-keto reductase gene from peach and its role in abiotic stress tolerance. *Phytochemistry* 104:30-36
- Kim, I.S., Y.S. Kim, and H.S. Yoon. 2012. Rice ASR1 protein with reactive oxygen species scavenging and chaperone-like activities enhances acquired tolerance to abiotic stresses in *Saccharomyces cerevisiae*. *Mol. Cells* 33:285-293.
- King, J.E. and D.J. Gifford. 1997. Amino acid utilization in seeds of loblolly pine during germination and early seedling growth: I. Arginine and arginase activity. *Plant Physiol.* 113:1125-1135.
- Kortstee, A.J., N. J. G. Appeldoorn, M.E.P. Oortwijn, and R.G.F. Visser. 2007. Differences in regulation of carbohydrate metabolism during early fruit development between domesticated tomato and two wild relatives. *Planta* 226:929-939.
- Krammer, G., W. Peter, S. Melanie, and S. Peter. 1991. Glycosidically bound aroma compounds in the fruits of *Prunus* species: apricot (*P. armeniaca*, L.), peach (*P. persica*, L.), yellow plum (*P. domestica*, L. ssp. *syriaca*). *J. Agric. Food Chem.* 39:778-781.
- Lara M.V., J. Borsani, C.O. Budde, M.A. Lauxmann, V.A. Lombardo, R. Murray, C.S. Andreo and M.F. Drincovich. 2009. Biochemical and proteomic analysis of 'Dixiland' peach fruit (*Prunus persica*) upon heat treatment. *J. Exp. Bot.* 60:4315-4333.
- Levi, A., A. Davis, A. Hernandez, P. Wechter, J. Thimmapuram, T. Trebitsh, Y. Tadmor, N. Katzir, V. Portnoy, and S. King. 2006. Genes expressed during the development and ripening of watermelon fruit. *Plant Cell Rep.* 25:1233-1245.
- Lichtenthaler H.K. 1987. Chlorophyll and carotenoids: Pigments of photosynthetic biomembranes. *Methods Enzymol.* 148:350-382.
- Lima, A.J.B., Á.A. Alvarenga, M.R. Malta, D. Gebert, and E.B. Lima. 2013. Chemical evaluation and effect of bagging new peach varieties introduced in southern Minas Gerais - Brazil. *Food Sci. Technol.* 33:434-440.
- Liu, J., Y. Sui, M. Wisniewskia, S. Drobycz, S.P. Tian, J. Norelli, and V. Hershkovitz. 2012. Effect of heat treatment on inhibition of *Monilinia fructicola* and induction of disease resistance in peach fruit. *Postharvest Biol. Technol.* 65:61-68.
- Liu, Y.L., F. Che, L.X. Wang, R. Meng, X.J. Zhang, and Z.Y. Zhao. 2013. Fruit coloration and anthocyanin biosynthesis after bag removal in non-red and red apples (*Malus × domestica* Borkh.). *Molecules* 18:1549-1563.
- Manrique-Trujillo, S.M., A.C. Ramirez-Lopez, E. Ibarra-Laciette, and M.A. Gomez-Lim. 2007. Identification of genes differentially expressed during ripening of banana. *J. Plant Physiol.* 164:1037-1050.
- Moing, A., L. Svanella, D. Rolin, M. Gaudillere, J.P. Gaudillere, and R. Monet. 1998. Compositional changes during the fruit development of two peach cultivars differing in juice acidity. *J. Am. Soc. Hortic. Sci.* 123:770-775.
- Nilo, R., C. Saffie, K. Lilley, R. Baeza-Yates, V. Cambiazo, R. Campos-Vargas, M. González, L.A. Meisel, J. Retamales, H. Silva, and A. Orellana. 2010. Proteomic analysis of peach fruit mesocarp softening and chilling injury using difference gel electrophoresis (DIGE). *BMC Genomics* 11:43.
- Polenta, G.A., J.J. Calvete, and C.B. González. 2007. Isolation and characterization of the main small heat shock protein induced in tomato pericarp by thermal treatment. *FEBS J* 274:6447-6455.
- Prinsi, B., A.S. Negri, C. Fedeli, S. Morgutti, N. Negrini, M. Cocucci, and L. Espen. 2011. Peach fruit ripening: A proteomic comparative analysis of the mesocarp of two cultivars with different flesh firmness at two ripening stages. *Phytochemistry* 72:1251-1262.
- Shen, J.Y., L.Wu, H.R. Liu, B. Zhang, X.R. Yin, Y.Q. Ge, and K.S. Chen. 2014. Bagging treatment influences production of C6 aldehydes and biosynthesis-related gene expression in peach fruit skin. *Molecules* 19:13461-13472.
- Shibata, H. and H. Ochiai. 1977. Purification and properties of δ -aminolevulinic acid dehydratase from radish cotyledons. *Plant Cell Physiol.* 18:421-429.
- Shulaev, V., S.S. Korban, B. Sosinski, A.G. Abbott, H.S. Aldwinckle, K.M. Folta, A. Iezzoni, D. Main, P. Arús, A.M. Dandekar, K. Lewers, S.K. Brown, T.M. Davis, S.E. Gardiner, D. Potter, and R.E. Veilleux. 2008. Multiple models for Rosaceae genomics. *Plant Physiol.* 147:985-1003.
- Sjögren, L.L.E., T.M. Stanne, B. Zheng, S. Sutinen, and A.K. Clarke. 2006. Structural and functional insights into the chloroplast ATP-dependent Clp protease in *Arabidopsis*. *The Plant Cell* 18:2635-2649.
- Watanabe, M., T. Oyamada, A. Suzuki, M. Murakami, S. Sugaya, S. Komori, and O. Arakawa. 2011. Comparison of sugar accumulation characteristics in 'Haruka' and 'Fuji' apple fruits. *Hortic. Res.* 10:565-571.
- Yan, S.C., J.Y. Chen, J.F. Kuang, W.X. Chen, and W.J. Lu. 2010. Expression of sHSP genes as affected by heat shock and cold acclimation in relation to chilling tolerance in plum fruit. *Postharvest Biol. Technol.* 55:91-96.
- Yan, S.P., Q.Y. Zhang, Z.C. Tang, W.A. Su, and W.N. Sun. 2006. Comparative proteomic analysis provides new insights into chilling stress responses in rice. *Mol. Cell Proteomics* 5:484-496.
- Yu, B., D. Zhang, C. Huang, M. Qian, X.Y. Zheng, Y.W. Teng, J. Su, and Q. Shu. 2012. Isolation of anthocyanin biosynthetic genes in red Chinese sand pear (*Pyrus pyrifolia* Nakai) and their expression as affected by organ/tissue, cultivar, bagging and fruit side. *Sci. Hortic.* 136:29-37.
- Zhang, B., J.Y. Shen, W.W. Wei, W.P. Xi, C.J. Xu, I. Ferguson, and K.S. Chen. 2010. Expression of genes associated with aroma formation derived from the fatty acid pathway during peach fruit ripening. *J. Agric. Food Chem.* 58: 6157-6165.
- Zhang, L., Z. Yu, L. Jiang, J. Jiang, H. Luo, and L. Fu. 2011. Effect of post-harvest heat treatment on proteome change of peach fruit during ripening. *J. Proteomics* 74:1135-1149.